

All Sites and Facilities
Standing Orders – Adults – NOT on Tolvaptan
Autosomal Dominant Polycystic Kidney Disease (ADKPD)

Page 2 of 2

Last Name:		
First Name (Preferred Name):		
Encounter number:	NH Number:	Chart Created: Y/N
Date of Birth:	Gender:	Age: Encounter Type:
Responsibility for Payment:		PHN:
Primary Care Physician/Attending Physician:		

PATIENT LABEL

Guidelines for Kidney Care Clinics:

Frequency of lab work in adults with Autosomal Dominant Polycystic Kidney Disease (not on Tolvaptan)

Extracted from BC Renal Guideline on Ordering, Reviewing and Follow-Up of Lab Work at www.bcrenalagency.ca

GFR (mL/min/1.73m ²)	≥ G3a ≥ 45	G3b 30-44	G4 15-29	G5 (<15) &/or Unstab
On ESA				
CBC, Retic Count (On ESA)	Monthly	Monthly	Monthly	Monthly
Ferritin, Serum Iron, TIBC, Iron Saturation (On ESA)	Q3 mos	Q3 mos	Q3 mos	Q3 mos
Not on ESA				
CBC (No ESA)	Q6 mos	Q3 mos	Q2 mos	Monthly
Ferritin, Serum Iron, TIBC, Iron Saturation (No ESA)	Q6 mos	Q3 mos	Q4 mos	Q3 mos
Other Tests				
Na ⁺ , K ⁺ , Cl ⁻ , Bicarbonate	Q6 mos	Q3 mos	Q2 mos	Monthly
Urea, Creatinine	Q6 mos	Q3 mos	Q2 mos	Monthly
Albumin, Ca ²⁺ , PO ₄	Q6 mos	Q3 mos	Q2 mos	Monthly
iPTH	Q12 mos	Q12 mos	Q6 mos	Q3 mos
AST, ALT, ALP, GGT, bilirubin, uric acid	Q6 mos	Q6 mos	Q4 mos	Q3 mos
Diabetes: Hgb A1c	Q6 mos	Q3 mos	Q4 mos	Q3 mos
Urine Albumin to Creatinine Ratio (ACR)	Q6 mos	Q3 mos	Q4 mos	Q3 mos
Urine Osmolality	Q6 mos	Q3 mos	Q4 mos	Q3 mos
24-hour urine for sodium, protein, creatinine, urea, osmolality. NOTE: Provide containers up to 6 L	Q12 mos	Q12 mos	Q12 mos	Q12 mos