

## **All Sites and Facilities**

## **Columbia Suicide Severity Rating Scale**

Page 1 of 1 PATIENT LABEL

Ask questions that are in bold and underlined.		Past month		
Ask questions 1 and 2		Yes	No	
1) Have you wished you were dead or wished you could go to sleep and not wake up?				
2) Have you had any actual thoughts of killing yourself?				
If yes to 2, ask questions 3, 4, 5 and 6. If no to 2, go directly to question 6.				
(e.g	ve you been thinking about how you might do this? I., "I thought about taking an overdose but I never made a specific plan as to en, where or how I would actually do it and I would never go through with it.")			
	ve you had these thoughts and had some intention of acting on them? opposed to "I have thoughts but I definitely will not do anything about them."			
	ve you started to work out or worked out the details of how you would kill urself? Do you intend to carry out this plan?			
6) Have you ever done anything, started to do anything, or prepared to do any thing to end your life?  Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.		Lifetime		
		Past 3 months		
If yes, ask: Was this within the past 3 months?				
Low Risk	Consider outpatient referral for community services			
Medium Risk	Itom 2 Develoring concult (Develoring Nursea/Social Morker) and consider nations eaterly proceditions			
High Risk	Item 4 and 5 - Psychiatric consultation and patient safety precautions Inpatient - Immediate notification of physician and/or psychiatric consultation	tation and patient safety precautions		
Medium Risk	Item 6 - Over 3 months ago. Psychiatric consult and consider patient safety	fety precautions		
High Risk	Item 6 three months ago or less: Immediate notification of physician and/or patient safety precautions	d/or psychiatric consultation and		
Level of risk identified:				
	Completed by:			
	· · · · · · · · · · · · · · · · · · ·			

